CONTENTS

	Page
Notes on the Composition of the Sorghum Plant. J. J. Willaman, R. M. West, D. O. Spriestersbach, and G. E. Holm....	1
Simple Method for Measuring the Acidity of Cereal Products: Its Application to Sulphured and Unsulphured Oats. Victor Birckner.....	33
Rate of Absorption of Soil Constituents at Successive Stages of Plant Growth. John S. Burd.....	51
Relation of the Concentration and Reaction of the Nutrient Medium to the Growth and Absorption of the Plant. D. R. Hoagland.....	73
Natural Carbonates of Calcium and Magnesium in Relation to the Chemical Composition, Bacterial Contents, and Crop-Producing Power of Two Very Acid Soils. S. D. Conner and H. A. Noyes.	119
Recent Studies on Sclerotium rolfsii Sacc. J. J. Taubenhaus....	127
Effect of Variation in Moisture Content on the Water-Extractable Matter of Soils. J. C. Martin and A. W. Christie.....	139
Pathology of Dourine with Special Reference to Microscopic Changes in Nerve Tissues and Other Structures. Robert J. Formad.....	145
Yellow-Berry in Hard Winter Wheat. Herbert F. Roberts...	155
Note on the European Corn Borer (Pyrausta nubilalis Hübner) and Its Nearest American Allies, with Description of Larvæ, Pupæ, and One New Species. Carl Heinrich.....	171
Bacterial Blight of Soybean. Florence M. Coerper.....	179
Do Mold Spores Contain Enzyms? Nicholas Kopeloff and Lillian Kopeloff.....	195
Nature and Control of Apple-Scald. Charles Brooks, J. S. Cooley, and D. F. Fisher.....	211
Nitrogen Metabolism of Two-Year-Old Steers. Sleeter Bull and H. S. Grindley.....	241
Development of the Pistillate Spikelet and Fertilization in Zea mays L. Edwin C. Miller.....	255
Response of Citrus Seedlings in Water Cultures to Salts and Organic Extracts. J. F. Breazeale.....	267
Physiological Study of the Parasitism of Pythium debaryanum Hesse on the Potato Tuber. Lon A. Hawkins and Rodney B. Harvey.....	275
Losses of Organic Matter in Making Brown and Black Alfalfa. C. O. Swanson, L. E. Call, and S. C. Salmon.....	299

Page

Cotton Rootrot Spots. C. S. SCOFIELD..... 305
Apple-Grain Aphis. A. C. BAKER and W. F. TURNER..... 311
Sunflower Silage. R. E. NEIDIG and LULU E. VANCE..... 325
Effect of Oxidation of Sulphur in Soils on the Solubility of Rock Phosphate and on Nitrification. O. M. SHEDD..... 329
Effect of Lime upon the Sodium-Chlorid Tolerance of Wheat Seedlings. J. A. LECLERC and J. F. BREAZEALE..... 347
Relation of Moisture in Solid Substrata to Physiological Salt Balance for Plants and to the Relative Plant-Producing Value of Various Salt Proportions. JOHN W. SHIVE..... 357
Treatment of Cereal Seeds by Dry Heat. D. ATANASOFF and A. G. JOHNSON..... 379
Meat Scraps versus Soybean Proteins as a Supplement to Corn for Growing Chicks. A. G. PHILIPS, R. H. CARR, and D. C. KENNARD..... 391
Further Studies of Sorosporella uvella, a Fungous Parasite of Noctuid Larvæ. A. T. SPEARE..... 399
Work and Parasitism of the Mediterranean Fruit Fly in Hawaii during 1918. H. F. WILLARD..... 441
Cyanogenesis in Sudan Grass: A Modification of the Francis-Connell Method of Determining Hydrocyanic Acid. PAUL MENAUL and C. T. DOWELL..... 447
European Frit Fly in North America. J. M. ALDRICH..... 451
Lepidoptera at Light Traps. W. B. TURNER..... 475
Life History of Eubiomyia calosomae, a Tachinid Parasite of Calosoma Beetles. C. W. COLLINS and CLIFFORD E. HOOD.... 483
Determination of Normal Temperatures by Means of the Equation of the Seasonal Temperature Variation and a Modified Thermograph Record. F. L. WEST, N. E. EDLEFSEN, and S. EWING..... 499
Temperature Relations of Certain Potato-Rot and Wilt-Producing Fungi. H. A. EDSON and M. SHAPOVALOV..... 511
Germination of Barley Pollen. STEPHEN ANTHONY and HARRY V. HARLAN..... 525
Invertase Activity of Mold Spores as Affected by Concentration and Amount of Inoculum. NICHOLAS KOPELOFF and S. BYALL..... 537
Basal Glumerot of Wheat. LUCIA MCCULLOCH..... 543
Effect of the Relative Length of Day and Night and Other Factors of the Environment on Growth and Reproduction in Plants. W. W. GARNER and H. A. ALLARD..... 553
Index..... 607

ERRATA AND AUTHORS' EMENDATIONS

Pages 60–63, legends to figures 3–8, "phosphate" should read "phosphorus."

Page 175, lines 45–46, Mr. Ainslie informs the author that he has never succeeded in rearing *Pyrausta ainsliei* to maturity on *Nelumbo lutea* as here stated. In the discussion of *P. ainsliei* it should be stated that the old name *P. obumbratalis* Lederer may eventually prove to apply to this species. Without an examination of the genitalia of Lederer's types in Europe, however, it can not be determined whether his name applies to this species or to *P. penitalis* Grote, if to either. In the meantime it is safest to employ a definite name to a definite concept.

Page 333, Table III, column 8, next to last line, and column 9, next to last line, "3.329" should read "3.320."

Page 335, Table VI, "phosphates" should read "phosphate."

Page 339, Table XIII, column 5, line 8, "92" should read "80."

Page 393, line 27, "in part by weight" should read "in parts by weight."

Page 394, Table I, column 9, line 10, "54.5" should read "5.45." Note *a*, "expressed in parts" should read "expressed in parts by weight."

Page 395, Table II, continued, column 6, line 4, "1 : 1.83" should read "1 : 0.83." Second part of table, column 6, lines 1 and 2, "1 : 1.05" and "1 : 1.37" should be transposed. Line 10 of text, "basal ratio" should read "basal ration."

Page 445, line 15 of text, "from 13.3 larvæ to each fruit in 1916 to 20.3 in 1917 and to 34.6 in 1918" should read "from 13.3 per cent in 1916 to 20.3 per cent in 1917 and to 34.6 per cent in 1918."

ILLUSTRATIONS

PLATES

NATURAL CARBONATES OF CALCIUM AND MAGNESIUM IN RELATION TO THE CHEMICAL COMPOSITION, BACTERIAL CONTENTS, AND CROP-PRODUCING POWER OF TWO VERY ACID SOILS

Page

PLATE 1. Pot cultures on acid yellow clay soil: A.—Wheat; B.—Red clover; C.—Red turnip beets. 126

PLATE 2. Pot cultures on acid black sand soil: A.—Wheat; B.—Red clover; C.—Red turnip beets. 126

RECENT STUDIES ON SCLEROTIUM ROLFSII SACC.

PLATE 3. *Sclerotium rolfsii:* A.—Two colonies in same plate, both apparently of the same sexual strain. B.—Large sclerotia from inoculated orange. C.—Late stage, showing cantaloupe reduced to a mass of mycelium and sclerotia. D.—Inoculated cantaloupe fruit, the point of infection being free and away from the bell jar glass. Earlier stage than C. E.—Infected cantaloupe, lying close to bell jar at point of infection. F.—Cross section of an infected cantaloupe, showing penetration of fungus into interior of fruit. 138

PLATE 4. *Sclerotium rolfsii:* A.—Cabbage artificially inoculated. B.—Cultures on artificial media. C.—Mustard-seed-like sclerotia on cantaloupe. D.—Formation of sclerotia at point of union of apparently plus and minus strains. 138

PLATE 5. *Sclerotium rolfsii:* A.—Sweet potato in seed bed, showing method of natural infection of young sprouts. B.—Infected Irish potato, "melter" stage. C.—Longitudinal section of infected sweet potato, showing nature of rot. 138

PLATE 6. *Sclerotium rolfsii:* A.—Dead tomato plants, showing fungus growing out on the surface of the soil in radial fans. B.—Growth on glass of bell jar, showing radial fans. 138

NOTE ON THE EUROPEAN CORN BORER (PYRAUSTA NUBILALIS HÜBNER) AND ITS NEAREST AMERICAN ALLIES, WITH DESCRIPTION OF LARVÆ, PUPÆ, AND ONE NEW SPECIES

PLATE 7. European corn borer and its nearest American allies: A.—Male genitalia of *Pyrausta nubilalis;* ventral view of organs spread; aedoeagus omitted. B.—Male genitalia of *P. nubilalis;* aedoeagus and penis. C.—Male genitalia of *P. ainsliei;* ventral view of organs spread; aedoeagus omitted. D.—Male genitalia of *P. penitalis;* ventral view of organs spread; aedoeagus omitted. E.—Forewing venation of *P. nubilalis;* male. F.—Hindwing venation of *P. nubialis;* male. 178

PLATE 8. Female genitalia of *Pyrausta* spp.: A.—Female genitalia of *P. nubilalis;* lateral view of organs. B.—Female genitalia of *P. nubilalis;* ventral view of plates posterior to genital opening. C.—Female genitalia of *P. penitalis;* ventral view of plates posterior to genital opening. D.—Female genitalia of *P. penitalis;* lateral view of organs. E.—Female genitalia of *P. ainsliei;* lateral view of organs. F.—Female genitalia of *P. ainsliei;* ventral view of plates surrounding genital opening 178

Page

PLATE 9. European corn borer and its nearest American allies: A.—Profile of cephalic end of pupa of *Pyrausta ainsliei*. B.—Profile of cephalic end of pupa of *P. penitalis*. C.—Profile of cephalic end of pupa of *P. nubilalis*. D.—Pupa of *P. nubilalis*, female; dorsal view. E.—Pupa of *P. nubilalis*, female; ventral view. F.—Caudal end of pupa of *P. nubilalis*, female; ventral view. G.—Caudal end of pupa of *P. penitalis*, female; ventral view..... 178

PLATE 10. European corn borer and its nearest American allies: A.—Dorsal view of head capsule; larva of *Pyrausta nubilalis*. B.—Lateral view of head capsule; larva of *P. nubilalis*. C.—Epipharynx; larva of *P. nubilalis*. D.—Labrum; larva of *P. nubilalis*. E.—Dorsal view of head capsule; larva of *P. penitalis*. F.—Lateral view of head capsule; larva of *P. penitalis*. G.—Dorsal view of head capsule; larva of *P. ainsliei*..... 178

PLATE 11. European corn borer and its nearset American allies: A.—Labium and maxillæ; larva of *Pyrausta nubilalis*. B.—Left maxilla; larva of *P. penitalis*. C.—Mandible; larva of *P. penitalis*. D.—Mandible; larva of *P. nubilalis*. E.—Character of skin granulations, highly magnified; larva of *P. nubilalis*. F.—Second thoracic and eighth and ninth abdominal segments of larva of *P. nubilalis*, showing granulose areas above seta VII. G.—Setal map of prothoracic, mesothoracic, third, eighth, and ninth abdominal segments; larva of *P. nubilalis*. H.—Crochet arrangement; abdominal proleg; larva of *P. nubilalis*..... 178

BACTERIAL BLIGHT OF SOYBEAN

PLATE A. Bacterial spot on leaves of soybean..... 194

PLATE 12. *Bacterium glycineum:* A.—Soybean petiole. Natural infection, showing exudate droplets. B.—Blighted soybean pod. Natural infection, showing young water-soaked lesions. C.—Blighted soybean pod. Natural infection, showing exudate droplets..... 194

PLATE 13. *Bacterium glycineum:* A.—Natural infection on soybean pods. B.—Artificial infection on soybean pods inoculated with type strain..... 194

PLATE 14. *Bacterium glycineum:* A.—Six-day-old potato agar slant culture, strain E. B and C.—Six-day-old potato agar slant cultures, type strain... 194

PLATE 15. *Bacterium glycineum:* A.—Colonies of petiole strain on potato agar plate. B.—Colonies of type strain on potato agar plate. C.—Colonies of strain E on potato agar plate. D.—Colony of seed strain on potato agar plate..... 194

PLATE 16. *Bacterium glycineum:* Soybean seedlings showing infection after inoculation in the greenhouse with type strain..... 194

PLATE 17. *Bacterium glycineum:* Blighted soybean leaves artificially inoculated in the field with type strain..... 194

PLATE 18. *Bacterium glycineum:* Soybean seedlings artificially inoculated in the greenhouse with type strain..... 194

DEVELOPMENT OF THE PISTILLATE SPIKELET AND FERTILIZATION IN ZEA MAYS L.

PLATE 19. Longitudinal section of the pistillate spikelet of corn at the time the silk is ready for pollination: r, rachilla; g, lower empty glume; g′, upper empty glume; l, lemma or flowering glume of the fertile flower; l′, lemma or flowering glume of the sterile flower; pa, palet of the fertile flower; pa′, palet of the sterile flower; sf, sterile flower; st, rudimentary stamen of the fertile flower; ova, ovary of the pistil; co, cavity of the ovary; sk, silk or style; sc, stylar canal; vbs, one of the fibro-vascular bundles of the silk.... 266

Page

PLATE 20. A.—Cross section of the tip of a very young cob. B.—Portion of a cross section of a young cob just back of the tip, showing the rudiment or primordium from which a pair of spikelets will develop. C.—Cross section of a rudiment at the beginning of its division into equal parts. D.—Cross section of a pair of spiklets in the process of development..... 266

PLATE 21. A.—Longitudinal section of the tip of a young cob. B.—Longitudinal section of the rudiment or primordium of a spikelet just back of the tip of a young cob: g, primordium of the lower empty glume. C.—Longitudinal section of the developing spikelet, showing the primordia of the lower and upper empty glumes: g, lower empty glume; g′, upper empty glume. D.—Longitudinal section of the developing spikelet at a little later stage than C: g, lower empty glume; g′, upper empty glume..... 266

PLATE 22. A.—Longitudinal section of the developing spikelet: g, lower empty glume; g′, upper empty glume; l, primordium of the lemma or flowering glume of the fertile flower; l′, primordium of the lemma or flowering glume of the sterile flower; sf, primordium of the sterile flower. B.—Longitudinal section of the developing spikelet: g and g′, empty glumes; l and l′, lemmas or flowering glumes; sf, primordium of the sterile flower; st, stamen of the fertile flower; p, palet of the fertile flower; pp, primordium of the pistil..... 266

PLATE 23. A.—Longitudinal section of the developing spikelet at the time the carpel or ovary wall has begun to develop: g and g′, empty glumes; l and l′, lemmas; sf, sterile flower; p and p′, palets; st, stamen; c and c′, rudiment of the carpel or ovary wall; c′ is the more rapidly growing part of the carpel. B.—Longitudinal section of a developing ovary: c and c′, developing ovary walls; c′ is the portion of the carpel from which the style or silk will develop; ov, ovule; ovc, primordium of the inner ovule coat..... 266

PLATE 24. A.—Longitudinal section of the fertile pistil at the time the silk has started to elongate: ovw, ovary wall; sc, stylar canal; ovc, ovule coats; ov, ovule; ms, megaspore mother cell; sk, silk. B.—Longitudinal section of the fertile pistil at the time the ovule has started to invert: ovw, ovary wall; sc, stylar canal; sk, silk; ovc, ovule coats; ms, megaspore-mother cell.. 266

PLATE 25. A.—Longitudinal section of the inverted ovule: ov, ovule; ovc, ovule coats; es, embryo sac; mic, micropyle; co, cavity of ovary. B.—Section through the stylar canal, showing its structure shortly before the silk emerges from the husk..... 266

PLATE 26. A.—Cross section of the megaspore mother cell. B.—Longitudinal section of the megaspore mother cell. C.—Longitudinal section of the 2-celled embryo sac..... 266

PLATE 27. A.—Longitudinal section of the 4-celled embryo sac. B.—Longitudinal section of an 8-celled embryo sac at the time the polar nuclei have started to migrate. C.—Longitudinal section of an 8-celled embryo sac after the polar nuclei have migrated..... 266

PLATE 28. Longitudinal section of a mature embryo sac just previous to fertilization: e, egg; sy, synergids; p, polars; ant, antipodals; pt, pollen tube; m, micropyle; v, vacuole; c, cytoplasm; ov, ovule coat..... 266

PLATE 29. A.—End of a silk: p, pollen grains; v, fibro-vascular bundles; h, hairs. B.—Tips of the hairs of the silk. C.—Section of a pollen grain showing the germ pore and the relative size of the vegetative and sperm nuclei. D.—Single hair of the silk, showing the general manner in which the pollen tube penetrates the sheath cells of the fibro-vascular bundle of the silk: p, pollen grain; h, hair; pt, pollen tube; sc, sheath cells of the fibro-vascu-

Page

lar bundle. E.—Longitudinal section of a fibro-vascular bundle of a silk, showing the position of the pollen tube as it grows down the silk: sc, sheath cells; x, xylem elements; p, parenchyma cells of the silk; pt, pollen tube, showing the enlarged tip..... 266

PLATE 30. A.—Cross section of a silk near its base, showing the position of the fibro-vascular bundles. B.—Cross section of a fibro-vascular bundle of the silk: x, xylem elements; sc, sheath cells; pt, pollen tube. C.—Longitudinal section through the sheath cells of the fibro-vascular bundles: sc, sheath cells; pc, parenchyma cells of the silk..... 266

PLATE 31. A.—Vegetative and sperm nuclei of the pollen grain: vn, vegetative nucleus; sn, sperm nuclei. B.—Longitudinal section of the lower portion of the embryo sac at the time of fertilization, reconstructed from two sections: pn, polar nuclei fusing; sn′, sperm nucleus fusing with a polar nucleus; e, egg; sn, sperm nucleus in the egg; pt, pollen tube; syn, synergid; v, vacuole..... 266

PLATE 32. A.—Longitudinal section of the embryo sac 12 hours after fertilization: end, endosperm nuclei; e, egg in which one of the daughter nuclei has already divided; ant, antipodals. B.—Longitudinal section of the embryo sac 36 hours after fertilization: end, endosperm; emb, embryo; ant, antipodal tissue. C.—Longitudinal section of the young embryo at the stage shown in B..... 226

RESPONSE OF CITRUS SEEDLINGS IN WATER CULTURES TO SALTS AND ORGANIC EXTRACTS

PLATE 33. A.—Grapefruit roots, 12 days old, grown in distilled water. B.—Grapefruit roots, 12 days old, grown in distilled water plus 100 parts per million water-soluble peat. C.—Lemon seedlings, 21 days old, grown in carbon-treated distilled water. D.—Lemon seedlings, 21 days old, grown in carbon-treated distilled water plus calcium carbonate. E.—Grapefruit seedlings, 20 days old, grown in carbon-treated water. F.—Grapefruit seedlings, 20 days old, grown in carbon-treated water plus calcium carbonate..... 274

PLATE 34. A.—Lemon seedlings, 16 days old, grown in distilled water as control. B.—Lemon seedlings, 16 days old, grown in 400 parts per million sodium carbonate. C.—Lemon seedlings, 16 days old, grown in 100 parts per million ammonia-soluble humus. D.—Lemon seedlings, 16 days old, grown in 100 parts per million ammonia-soluble humus plus 400 parts per million sodium carbonate. E.—Citrus seedlings grown in carbon-treated water. F.—Citrus seedlings grown in 4/10 per cent sodium nitrate ($NaNO_3$). G.—Citrus seedlings grown in 4/10 per cent sodium nitrate with calcium carbonate ($CaCO_3$)..... 274

PHYSIOLOGICAL STUDY OF THE PARASITISM OF PYTHIUM DEBARYANUM HESSE ON THE POTATO TUBER

PLATE 35. A.—Apparatus used in determining the pressure required to puncture the tissue of potato tubers. B.—Glass rod with attached needle..... 298

PLATE 36. Photographs enlarged from portions of a motion photomicrograph, showing the method of cell wall penetration by Pythium hyphae. A.—Shows hypha growing toward the potato cell wall. B.—Shows hypha attached to wall and about to penetrate. C.—The tip has just broken through the wall. D.—The penetration is complete..... 298

Page

PLATE 37. Photographs enlarged from portions of a motion photomicrograph, showing the method of cell wall penetration by Pythium hyphae. A.—Shows the hypha growing against the potato cell wall. B.—A little later stage than A. C.—The tip has broken through as a small tube. D.—Penetration is complete 298

EFFECT OF LIME UPON THE SODIUM-CHLORID TOLERANCE OF WHEAT SEEDLINGS

PLATE 38. A.—Seedlings grown in (1) distilled water, (2) 500 parts per million sodium-chlorid solution, (3) 1,000 parts per million sodium-chlorid solution, (4) 2,000 parts per million sodium-chlorid solution, (5) 3,000 parts per million sodium-chlorid solution, and (6) 4,000 parts per million sodium-chlorid solution. B.—Seedlings grown in sand and watered with (1) distilled water, (2) 500 parts per million sodium-chlorid solution, (3) 1,000 parts per million sodium-chlorid solution, (4) 2,000 parts per million sodium-chlorid solution, (5) 3,000 parts per million sodium-chlorid solution, and (6) 4,000 parts per million sodium-chlorid solution 356

PLATE 39. Seedlings grown in water, sand, and clay, showing effects of (1) distilled water, (2) 500 parts per million sodium-chlorid solution, (3) 1,000 parts per million sodium-chlorid solution, (4) 1,500 parts per million sodium-chlorid solution, (5) 2,000 parts per million sodium-chlorid solution, (6) 3,000 parts per million sodium-chlorid solution, and (7) 4,000 parts per million sodium-chlorid solution 356

PLATE 40. A.—Seedling, 9 days old, grown in clay watered with (1) distilled water, (2) 500 parts per million sodium-chlorid solution, (3) 1,000 parts per million sodium-chlorid solution, (4) 1,500 parts per million sodium-chlorid solution, (5) 2,000 parts per million sodium-chlorid solution, (6) 3,000 parts per million sodium-chlorid solution, and (7) 4,000 parts per million sodium-chlorid solution. B.—Seedlings, 3 days old, removed from (1) 4,000 parts per million sodium-chlorid solution, (2) sand watered with 4,000 parts per million sodium-chlorid solution, and (3) clay watered with 4,000 parts per million sodium-chlorid solution 356

PLATE 41. A.—Seedlings grown in sand and in the following solutions filtered through sand: (1) distilled water, (2) 1,000 parts per million of sodium chlorid, (3) 1,500 parts per million of sodium chlorid, (4) 2,000 parts per million of sodium chlorid, (5) 3,000 parts per million of sodium chlorid, and (6) 4,000 parts per million of sodium chlorid. B.—Seedlings grown in clay and in the following solutions filtered through clay: (1) distilled water, (2) 1,000 parts per million of sodium chlorid, (3) 1,500 parts per million of sodium chlorid, (4) 2,000 parts per million of sodium chlorid, (5) 3,000 parts per million of sodium chlorid, and (6) 4,000 parts per million of sodium chlorid 356

PLATE 42. A.—Seedlings, 10 days old, removed from sand watered with (1) distilled water, (2) 1,000 parts per million sodium-chlorid solution, (3) 1,500 parts per million sodium-chlorid solution, (4) 2,000 parts per million sodium-chlorid solution, (5) 3,000 parts per million sodium-chlorid solution, and (6) 4,000 parts per million sodium-chlorid solution. B.—Seedlings, 10 days old, removed from clay watered with (1) distilled water, (2) 1,000 parts per million sodium-chlorid solution, (3) 1,500 parts per million sodium-chlorid solution, (4) 2,000 parts per million sodium-chlorid solution, (5) 3,000 parts per million sodium-chlorid solution, and (6) 4,000 parts per million sodium-chlorid solution 356

Page

PLATE 43. A.—Seedlings grown in (1) 4,000 parts per million sodium-chlorid solution, (2) 6,000 parts per million sodium-chlorid solution filtered through soil, (3) 8,000 parts per million sodium-chlorid solution filtered through soil, (4) 4,000 parts per million sodium-chlorid solution filtered through soil, (5) 4,000 parts per million sodium-chlorid solution added to distilled water that had previously been filtered through soil, and (6) 4,000 parts per million sodium-chlorid solution+100 parts per million each of sodium nitrate, potassium chlorid, and sodium phosphate. B.—Seedlings grown in (1) 4,000 parts per million sodium-chlorid solution, (2) 4,000 parts per million sodium-chlorid solution+30 parts per million of calcium sulphate, (3) 4,000 parts per million sodium-chlorid solution+30 parts per million of calcium oxid, and (4) 4,000 parts per million sodium-chlorid solution+30 parts per million of magnesium bicarbonate 356

PLATE 44. A.—Seedlings grown in (1) distilled water, (2) 4,000 parts per million sodium-chlorid solution + 30 parts per million of calcium sulphate, (3) 4,000 parts per million sodium-chlorid solution + 30 parts per million of calcium oxid, (4) 4,000 parts per million sodium-chlorid solution + 30 parts per million of magnesium sulphate, and (5) 4,000 parts per million sodium-chlorid solution + 30 parts per million of barium chlorid. B.—Seedlings grown in (1) distilled water, (2) 4,000 parts per million sodium-chlorid solution + 30 parts per million of potassium chlorid, (3) 4,000 parts per million sodium-chlorid solution + 30 parts per million of sodium nitrate, (4) 4,000 parts per million sodium-chlorid solution + 30 parts per million of sodium phosphate, (5) 4,000 parts per million sodium-chlorid solution + 30 parts per million of ferric chlorid, and (6) 4,000 parts per million sodium-chlorid solution + 30 parts per million of potassium alum. 356

PLATE 45. A.—Seedlings grown in sand watered with (1) 4,000 parts per million sodium-chlorid solution and (2) 4,000 parts per million sodium-chlorid solution + 30 parts per million of calcium sulphate. B.—Seedlings grown in (1) 4,000 parts per million sodium-sulphate solution, (2) 4,000 parts per million sodium-sulphate solution + 30 parts per million of calcium sulphate, (3) 4,000 parts per million sodium-sulphate solution + 30 parts per million of calcium oxid, (4) 4,000 parts per million sodium-sulphate solution + 30 parts per million of magnesium sulphate, and (5) 4,000 parts per million sodium-sulphate solution + 30 parts per million of barium chlorid 356

PLATE 46. A.—Seedlings, 3 days old, grown in (1) distilled water, (2) 4,000 parts per million sodium-sulphate solution, (3) 4,000 parts per million sodium-sulphate solution + 30 parts per million of potassium chlorid, (4) 4,000 parts per million sodium-sulphate solution + 30 parts per million of sodium nitrate, (5) 4,000 parts per million sodium-sulphate solution + excess ferric hydrate, and (6) 4,000 parts per million sodium-sulphate solution + aluminum hydrate. B.—Seedlings, 11 days old, grown in (1) 2,500 parts per million sodium-bicarbonate solution, (2) 2,500 parts per million sodium-bicarbonate solution + 30 parts per million of sodium nitrate, (3) 2,500 parts per million sodium-bicarbonate solution + 30 parts per million of potassium chlorid, (4) 2,500 parts per million sodium-bicarbonate solution + 30 parts per million of magnesium sulphate, and (5) 2,500 parts per million sodium-bicarbonate solution + 30 parts per million of calcium oxid 356

PLATE 47. A.—Seedlings grown in (1) 4,000 parts per million sodium-chlorid solution, (2) distilled water, (3) 4,000 parts per million sodium-chlorid solution + 40 parts per million of calcium oxid, (4) 4,000 parts per million

Page

sodium-chlorid solution + 30 parts per million of calcium oxid, and (5) 4,000 parts per million sodium-chlorid solution + 20 parts per million of calcium oxid. B.—Seedlings grown in (1) 4,000 parts per million sodium-chlorid solution, (2) 4,000 parts per million sodium-chlorid solution + 15 parts per million of calcium oxid, (3) 4,000 parts per million sodium-chlorid solution + 10 parts per million of calcium oxid, (4) 4,000 parts per million sodium-chlorid solution + 2 parts per million of calcium oxid, and (5) 4,000 parts per million sodium-chlorid solution + 1 part per million of calcium oxid..... 356

TREATMENT OF CEREAL SEEDS BY DRY HEAT

PLATE 48. Chevalier barley plants from untreated and treated seed referred to in Table III: A.—Plants from untreated seed. B.—Plants from seed treated with dry heat..... 390

PLATE 49. A.—Basal portions of 10 representative Chevalier barley plants, from untreated and treated seed, selected from the lots illustrated in Plate 48. B.—Representative Kubanka wheat seedlings from untreated seed (6 plants at left) and treated seed (5 plants at right) referred to on p. 385-386..... 390

MEAT SCRAPS VERSUS SOYBEAN PROTEINS AS A SUPPLEMENT TO CORN FOR GROWING CHICKS

PLATE 50. A.—Cockerel No. 43, lot 13, fed on basal ration only. B.—Pullet No. 44, lot 13, fed on basal ration only. C.—Cockerel No. 54, lot 10, fed on basal ration plus 5 parts meat scraps protein and 5 parts soybean meal protein. D.—Pullet No. 82, lot 6, fed on basal ration plus 10 parts soybean meal protein..... 398

FURTHER STUDIES OF SOROSPORELLA UVELLA, A FUNGOUS PARASITE OF NOCTUID LARVÆ

PLATE 51. *Sorosporella uvella:* A.—Infected cutworm torn open, exposing the resting-spore aggregations. B.—A single resting-spore aggregation. C.—A portion of a resting-spore aggregation germinating in water, showing promycelial-like germination, and conidia. D.—A colony of young resting spores from an infected insect, showing the manner in which they reproduce. E.—Isolated mature resting spores, some of which show the ruptured walls of previously cohering spores. F.—Mature resting spores germinating in water. G.—Isaria-like fascicle of cohering conidiiferous hyphæ which developed when the resting spores of an infected larva were allowed to germinate in a moist chamber. H.—Portion of a mature resting-spore aggregation. I.—Conidia, or secondary spores. J.—Conidia, or secondary spores, germinating. K.—Portion of a section through the body of an infected cutworm which had been placed in a moist chamber to induce germination of the resting spores, showing the usual type of conidiophores. L.—Mature resting spore germinating in water, showing conidiophore with verticillately arranged sterigmata. M.—Mature resting spores germinating on nutrient agar, showing sessile sterigmata and conidia at one place and young resting spores arising by budding at another place. N.—Early stages in the germination of mature resting spores in water. O.—Advanced stages in the germination of resting spores in water. P.—Sterigmata, showing method of conidial abjunction. Q.—Enlarged view of conidial germination. R.—Enlarged view of conidia. S.—Torula-like reproduction in culture..... 440

Page

PLATE 52. *Sorosporella uvella:* A.—Portion of a fat body from an infected insect which is being destroyed by an adhering group of young resting spores. B, C.—Phagocytes from an infected cutworm with blastocysts of the fungus imbedded in them. D, E.—Phagocytes distorted in form and partially disintegrated as a result of the action of the enclosed blastocysts. F.—The terminal portion of a complexly branched hypha from nutrient agar, showing sterigmata and conidia. G.—Normal blood corpuscles of *Feltia jaculifera*. H.—Resting spore germinating on nutrient agar, showing hyphæ, some of which are producing resting spores. I.—Blastocysts, showing method of reproduction. J.—An aggregation of cohering leucocytes in the substance of some of which blastocyets are to be seen. K.—Portion of the intestine of an infected *Feltia jaculifera*, showing longitudinal and transverse muscles connected by a frail membrane within the folds of which blastocysts may be seen. L.—Colony of young resting spores from a culture which are giving rise to hyphæ. M.—Colony from a Petri dish culture, showing a mass of young resting spores, some of which have germinated in situ by giving rise directly to sterigmata and conidia, and others that are producing hyphæ. N.—Small phagocytic complex. O.—Portion of one hypha rising from a resting spore that had germinated in a Petri dish culture, showing sterigmata and conidia..... 440

PLATE 53. *Sorosporella uvella:* A. Photomicrograph of a stained blood smear from an infected cutworm, showing a number of free-floating blastocysts and one leucocyte within which are to be seen two blastocysts. B.—Photomicrograph of a water mount of blastocysts which developed on Uschinsky's medium after inoculation..... 440

PLATE 54. *Sorosporella uvella:* A.—General appearance of the thallus when grown on Molisch's medium, showing brainlike convolutions and crateriform structure. B.—Streak culture upon beerwort agar, showing shiny gelatinous-appearing, grapelike bunches of mature resting spores, some of which are germinating..... 440

PLATE 55. *Sorosporella uvella:* A.—Infected larva of *Feltia* sp. torn open, showing Isaria-like fascicles of conidiiferous hyphæ that sometimes develop when the resting spores are allowed to germinate in a moist chamber. B.—Photomicrograph of a stained cross section of the heart, within the cavity of which numerous blastocysts are to be seen..... 440

PLATE 56. *Sorosporella uvella:* A, B.—Photomicrographs of phagocytic complexes, showing blastocysts incorporated in the substance of the phagocytes.. 449

EUROPEAN FRIT FLY IN NORTH AMERICA

PLATE 57. *Oscinis frit:* A.—Adult. B.—Puparium..... 474

LIFE HISTORY OF EUBIOMYIA CALOSOMAE, A TACHINID PARASITE OF CALOSOMA BEETLES

PLATE 58. *Eubiomyia calosomae:* A.—Egg, showing flat side and exit hole of larva. B.—Mouth hook of third-stage larva. C.—Third-stage larva after preservation in alcohol. D.—Adult of *Calosoma sycophanta* containing two second-stage or hibernating larvae of Eubiomyia..... 498

PLATE 59. *Eubiomyia calosomae:* A.—Puparium. B.—Anal stigmata of puparium from slide mount. C.—Section of spine area on segment of puparium. D.—Adult female..... 498

GERMINATION OF BARLEY POLLEN

Page

PLATE 60. A.—Normal pollen. B.—Bulging of the intine through imbibition. C.—The effect of free water when the intine is not ruptured. D.—Pollen germinating on a stigma hair; the pollen grain loosened so as to show the tube. E.—Germinating pollen. F, G.—Pollen tube..... 536

PLATE 61. A.—Normal pollen grains. B.—Same pollen grains after an exposure of two minutes to dry air..... 536

BASAL GLUMEROT OF WHEAT

PLATE 62. Basal glumerot of wheat: A.—Heads showing diseased glumes. B.—Glumes diseased at base. C.—Grains black and full of bacteria at base. D.—Young wheat leaves five days after inoculation with *Bacterium atrofaciens*. E.—Glumes inoculated with strain 478. F.—Glumes inoculated with strain 399. *a*, Inner face..... 552

PLATE 63. Various types of colonies of *Bacterium atrofaciens:* A.—Moderately crowded colonies on agar plate, 3 days old. B.—Well-isolated colonies on agar plate, 2 days old. C.—Single colony on agar plate, 20 days old. D.—Well-isolated colonies on agar plate, 3 days old. E.—Crowded colonies on agar plate, 2 days old. A and E represent the ordinary type of young colony. F.—Flagella (Van Ermengem's stain)..... 552

EFFECT OF THE RELATIVE LENGTH OF DAY AND NIGHT AND OTHER FACTORS OF THE ENVIRONMENT ON GROWTH AND REPRODUCTION IN PLANTS

PLATE 64. A.—Small dark chamber used in the 1918 experiments. B.—Larger dark house used in the 1919 tests..... 606

PLATE 65. A.—Peking soybeans exposed to the light for 7 hours daily. B.—Control planting of Peking soybeans exposed to the light for the whole day—that is, kept out of doors day and night..... 606

PLATE 66. A.—Peking soybeans exposed to the light for 7½ hours daily, beginning with the blossoming period. B.—Control planting of Peking soybeans kept out of doors throughout the test..... 606

PLATE 67. A.—Biloxi soybeans exposed to the light for 7 hours daily. B.—Control planting of Biloxi soybeans kept out of doors during the test..... 606

PLATE 68. Biloxi soybeans: A.—Plants in box on left exposed to the light for 5 hours daily. B.—Plants in box on left exposed to light daily for 7 hours; those on right exposed 12 hours daily..... 606

PLATE 69. A.—Biloxi soybeans. Plants in box on right exposed to light from daylight to 10 a. m. and from 2 p. m. to dark, a total of 9 to 10 hours daily. Plants in box on left exposed to light from 6 a. m. to 6 p. m., or 12 hours daily. B.—Radish plant in which the seed stalk was transformed into a vegetative shoot through the influence of the decreasing length of day.

PLATE 70. A.—Maryland Mammoth tobacco in 8-inch pots exposed to light from 9 a. m. to 4 p. m. daily. B.—Control series of Maryland Mammoth plants kept out of doors..... 606

PLATE 71. A.—Maryland Mammoth tobacco in 12-quart buckets exposed to light from 6 a. m. to 6 p. m., or 12 hours daily. B.—Maryland Mammoth tobacco in 12-quart buckets exposed to light from 9 a. m. to 4 p. m., or 7 hours daily..... 606

PLATE 72. A.—Control series of Maryland Mammoth tobacco in 12-quart buckets left out of doors during the experiment. B.—*Aster lunariifolius* L. Plants in box on left exposed to light from 9 a. m. to 4 p. m. daily. Plants in box on right left out of doors during the test..... 606

Page

PLATE 73. A.—*Phaseolus vulgaris* from Peru and Bolivia exposed to light from 9 a. m. to 4 p. m. B.—Control series of Phaseolus left out of doors during the experiment 606

PLATE 74. A.—*Mikania scandens* L. exposed to light from 9 a. m to 4 p. m. daily. B.—Control plants of Mikania left out of doors during the test 606

PLATE 75. A.—Ragweed. Plants on left exposed to light from 9 a. m. to 4 p. m. daily. Plants on right left out of doors as controls. B.—Radish. Plants in box on left exposed to light from 9 a. m. to 4 p. m. daily 606

PLATE 76. A.—Portion of stem of Maryland Mammoth tobacco plant, showing the sharp delimitation of dying back of the original growth as controlled by the appearance of new shoots. B.—Triangular type of shade with cheesecloth covering, used in the 1916 test with soybeans 606

PLATE 77. A.—Shade cloth, 6 by 6 mesh. B.—Shade cloth, 8 by 10 mesh. C.—Shade cloth, 12 by 12 mesh. D.—Shade cloth, 12 by 20 mesh. E.—Standard cheesecloth used in 1916 experiments 606

PLATE 78. A.—Soybeans growing in box set in soil (covers removed), shaded with 12 by 12 mesh netting. B.—Series of consecutive plantings of soybeans in the field during the summer 606

PLATE 79. A.—Soybeans growing in four boxes set in the soil and provided with removable covers. B.—Carrots. Plants in box on left exposed to light from 9 a. m. to 4 p. m. daily 606

TEXT FIGURES

NOTES ON THE COMPOSITION OF THE SORGHUM PLANT

FIG. 1. Development of the proximate and mineral constituents of the sorghum plant in the later stages of growth 4

2. Total weight of proximate constituents in a single sorghum plant during the later stages of growth 6

3. Development of the proximate and mineral constituents of the leaves of sorghum 7

4. Total weight of proximate constituents in the leaves of a single sorghum plant 7

5. Development of the proximate and mineral constituents of the seed heads of sorghum 8

6. Total weight of proximate constituents in the seed head of a single sorghum plant 9

7. Development of the proximate and mineral constituents of the bagasse of sorghum 10

8. Total weight of proximate constituents in the bagasse of a single sorghum plant 11

9. Development of the percentages of dry matter in the various parts of the sorghum plant 12

10. Development of the crude protein in the various parts of the sorghum plant 13

11. Development of the ether extract in the various parts of the sorghum plant 14

12. Development of the crude fiber in the various parts of the sorghum plant 15

13. Development of the nitrogen-free extract in the various parts of the sorghum plant 16

14. Development of the ash content in the various parts of the sorghum plant 17

Page

FIG. 15. Development of the composition of the ash in various parts of the sorghum plant 18
16. Distribution of the sugars and other solids in the various joints of sorghum cane 24
17. Development of the constituents of the juice of sorghum 25
18. Comparison of Early Amber sorghum grown in Minnesota with that grown in the District of Columbia 26

RATE OF ABSORPTION OF SOIL CONSTITUENTS AT SUCCESSIVE STAGES OF PLANT GROWTH

FIG. 1. Growth reached by barley at different periods of cutting. Experiment of 1910 59
2. Growth reached by barley at different periods of cutting. Experiment of 1917 59
3. Relation of growth of barley to absorption of potassium, nitrogen, phosphate, calcium, and magnesium, for entire plant except roots. Experiment of 1916 60
4. Relation of growth of barley to absorption of potassium, nitrogen, phosphate, calcium, and magnesium, for entire plant except roots. Experiment of 1917 61
5. Relation of growth of barley to absorption of potassium, nitrogen, phosphate, calcium, and magnesium, for stems and leaves. Experiment of 1916 62
6. Relation of growth of barley to absorption of potassium, nitrogen, phosphate, calcium, and magnesium, for heads. Experiment of 1916 62
7. Relation of growth of barley to absorption of potassium, nitrogen, phosphate, calcium, and magnesium, for stems and leaves. Experiment of 1917 63
8. Relation of growth of barley to absorption of potassium, nitrogen, phosphate, calcium, and magnesium, for heads. Experiments of 1917 63
9. Absorption of nitrogen by barley, expressed as the nitrate (NO_3) equivalent and computed to parts per million of soil 66
10. Absorption of potassium by barley, computed to parts per million of soil 67
11. Absorption of calcium by barley, computed to parts per million of soil 67
12. Absorption of magnesium by barley, computed to parts per million of soil 68
13. Absorption of phosphorus by barley, expressed as the phosphate (PO_4) equivalent and computed to parts per million of soil 68

RELATION OF THE CONCENTRATION AND REACTION OF THE NUTRIENT MEDIUM TO THE GROWTH AND ABSORPTION OF THE PLANT

FIG. 1. Water cultures, series 1. Graph showing net absorption in parts per million from solutions of four concentrations: Solution I with concentration of 0.10 atmospheres. Solution II with concentration of 0.32 atmospheres. Solution III with concentration of 0.85 atmospheres. Solution IV with concentration of 2.07 atmospheres 88

Page

FIG. 2. Water cultures, series 1. Graphs showing comparison between percentage of total nutrient absorbed and percentage of total water transpired with solutions of four concentrations 90
3. Water cultures, series 2. Graph showing absorption of nutrients from solutions of three concentrations during latter part of growth cycle. 92
4. Water cultures, series 3. Graph showing absorption of nutrients from solutions of two concentrations and two reactions 94

RECENT STUDIES ON SCLEROTIUM ROLFSII SACC.

FIG 1. A, intracellular nature of *Sclerotium rolfsii* hyphae in cantaloupe tissue; B and C, manner in which the fungus pierces host cells; D, E, and F, method of budding and formation of new mycelial growth; G, manner of growth of mycelium, forming strands; H, dissolved middle lamellae of host cells 134

YELLOW-BERRY IN HARD WINTER WHEAT

FIG. 1. Percentage by weight of yellow-berry kernels in pure-bred wheats and their control, 1908 162
2. Relation between date of harvesting and percentage of yellow-berry .. 163

BACTERIAL BLIGHT OF SOYBEAN

FIG. 1. *Bacterium glycineum:* From 72-hour growth on potato agar, stained by Duckwall's method to show flagella 182

NATURE AND CONTROL OF APPLE-SCALD

FIG. 1. Relative rate of cooling of apples in commercial and ventilated barrels: A, Grimes apples in a storage room already filled with cold fruit; B, York Imperial apples in a storage room still receiving a large bulk of warm fruit 232
2. Relative carbon-dioxid content of the air in ventilated and commercial barrels during the first weeks of storage: A, Grimes apples in a storage room already filled with cooled fruit; B, York Imperial apples in a storage room still receiving a large bulk of fresh fruit 233

NITROGEN METABOLISM OF TWO-YEAR-OLD STEERS

FIG. 1. Nitrogen metabolism of steers in the maintenance lot 243
2. Nitrogen metabolism of steers in the one-third feed lot 249
3. Nitrogen metabolism of steers in the two-thirds feed lot 250
4. Nitrogen metabolism of steers in the full feed lot 251

PHYSIOLOGICAL STUDY OF THE PARASITISM OF PYTHIUM DEBARYANUM HESSE ON THE POTATO TUBER

FIG. 1. Drawing to illustrate growth of a Pythium hypha in potato tissue 290
2. Drawing to illustrate method of cell-wall penetration in cells I, IV, and V 291

COTTON ROOTROT SPOTS

FIG. 1. Diagram of plot B5–4, showing by the brushlike lines the portions of the rows in which the cotton plants were killed by rootrot in 1916 306
2. Diagram of plot B5–4, showing by the heavy lines the portions of the rows in which the cotton plants were killed by rootrot in 1917 307

Page

FIG. 3. Diagram of plot B5–4, showing by the diagonal hatching the portions of the rows in which the cotton plants were killed by rootrot in 1918. 307
4. Diagram of plot B5–4, showing the portions of the rows in which the cotton plants were killed each year by rootrot in 1916, 1917, and 1918. 308
5. Diagram of plot A4–19, showing the portions of the rows in which the cotton plants were killed each year by rootrot in 1916, 1917, and 1918. 308
6. Diagram of plot A6–3, showing the portions of the rows in which the cotton plants were killed each year by rootrot in 1916, 1917, and 1918. 309
7. Diagram of plot B5–3, showing the portions of the rows in which the cotton plants were killed each year by rootrot in 1916, 1917, and 1918. 309

APPLE-GRAIN APHIS

FIG. 1. Effect of temperature upon the duration of the larval stages of the apple-grain aphis. 318

RELATION OF MOISTURE IN SOLID SUBSTRATA TO PHYSIOLOGICAL SALT BALANCE FOR PLANTS AND TO THE RELATIVE PLANT-PRODUCING VALUE OF VARIOUS SALT PROPORTIONS

FIG. 1. Diagrams showing the position of the cultures producing the nine highest yields of wheat tops in each series. 364
2. Average absolute yields of wheat tops for low, medium, and high moisture content of sand cultures. 366
3. Diagrams showing the position of the cultures producing the nine highest yields of wheat roots in each series. 368
4. Average absolute yields of wheat roots for low, medium, and high moisture content of sand cultures. 370
5. Amounts of water lost by transpiration from wheat plants grown in sand cultures with low, medium, and high moisture content. 372
6. Water requirement (in cubic centimeters per gram) of wheat tops grown in sand cultures with low, medium, and high moisture content. 374
7. Water requirement (in cubic centimeters per gram) of wheat roots grown in sand cultures with low, medium, and high moisture content. 374

MEAT SCRAPS VERSUS SOYBEAN PROTEINS AS A SUPPLEMENT TO CORN FOR GROWING CHICKS

FIG. 1. Graph showing the rate of growth of males and females in all lots. 392

EUROPEAN FRIT FLY IN NORTH AMERICA

FIG. 1. Map showing distribution of *Oscinis frit*. 454
2. Egg of *Oscinis frit*. 455
3. Newly hatched larva of *Oscinis frit*. 455
4. Full-grown larva of *Oscinis frit*. 456
5. *Oscinis frit:* Mouth hooks of full-grown larva. 456
6. *Oscinis frit:* Female abdomen, distended with eggs and ovipositor protruded. 457
7. *Oscinis frit:* Male genitalia, highly magnified. 457

LEPIDOPTERA AT LIGHT TRAPS

FIG. 1. Light traps: A, opening for cone; BBBB, plates of glass; C, glass tube to convey acid; D, glass jar to hold cyanid; E, glass funnel. 476

DETERMINATION OF NORMAL TEMPERATURES BY MEANS OF THE EQUATION OF THE SEASONAL TEMPERATURE VARIATION AND A MODIFIED THERMOGRAPH RECORD

Page

FIG. 1. Synoptic chart of annual temperature marches at selected stations in the United States 501
2. Mean monthly temperatures for Utah 502
3. Average Utah thermograph records for various seasons 504
4. Actual thermograph records taken at widely separated stations in the United States 505
5. Hourly variation of temperatures expressed in percentages of the mean. 507
6. Effect of storm on diurnal variation of temperature 509

TEMPERATURE RELATIONS OF CERTAIN POTATO-ROT AND WILT-PRODUCING FUNGI

FIG. 1. Graph showing the rate of growth of *Fusarium coeruleum* on potato agar at different temperatures 513
2. Graph showing the rate of growth of *Fusarium discolor* var. *sulphureum* on potato agar at different temperatures 514
3. Graph showing the rate of growth of *Fusarium eumartii* on potato agar at different temperatures 515
4. Graph showing the rate of growth of *Fusarium oxysporum* on potato agar at different temperatures 516
5. Graph showing the rate of growth of *Fusarium radicicola* on potato agar at different temperatures 517
6. Graph showing the rate of growth of *Fusarium trichothecioides* on potato agar at different temperatures 518
7. Graph showing the rate of growth of *Verticillium albo-atrum* No. 426 on potato agar at different temperatures 519
8. Graph showing the rate of growth of *Verticillium albo-atrum* No. 427 on potato agar at different temperatures 519
9. Graph showing the total amounts of growth produced by different potato-rot and wilt-fungi during the first seven days on potato agar at different temperatures 520

GERMINATION OF BARLEY POLLEN

FIG. 1. Period of receptivity of the stigma under field conditions as shown by the number of seed produced when 40 flowers were pollinated on emasculation and the same number on six successive days after emasculation 528
2. Number and percentage of seed produced by the use of pollen from anthers at various stages of development before and after dehiscence.. 529

EFFECT OF THE RELATIVE LENGTH OF DAY AND NIGHT AND OTHER FACTORS OF THE ENVIRONMENT ON GROWTH AND REPRODUCTION IN PLANTS

FIG. 1. Graph showing the shortening of the vegetative period preceding flowering soybeans which results from progressively later planting during the growing season 568
2. Graph showing changes in length of day during the growing season in the latitude of Washington, D. C. 569
3. Graph showing the progressive decrease in height attained by Biloxi soybeans as the date of planting is delayed beyond late spring 570